



GLOW BLEND

GHK-cu 50mg: 3ml BAC

BPC-157 10mg: 2ml BAC

TB 500 10mg: 1ml BAC

GHK-cu

BPC-157

TB-500

12 Units Daily

12 Units Daily

20 Units 3x Week

FOR OPTIMAL RESULTS:

(Continually Repeat 4-6 Weeks On, 4 Weeks Off)




www.precisionpeptideco.com

code: PEP5

GLOW BLEND

STEP 2

Example to visually shows how much is drawn up into an insulin syringe.





Reconstitute Peptide + BAC Water

GHK-cu 50mg + 3ml BAC


BPC-157 10mg + 2ml BAC

TB-500 10mg + 1ml BAC

STEP 1



STEP 3



FOR OPTIMAL RESULTS: Continually Repeat 4-6 Weeks On, 4 Weeks Off

www.precisionpeptideco.com ~ code: PEP5

Custom Glow Peptide Blend **For Anti-Aging, Skin Glow, and Repair**

Peptide Supplies:

- GHK-Cu: 50 mg (1 vial)
- BPC-157: 10 mg (1 vial)
- TB-500: 10 mg (1 vial)
- BAC Water:- 6mL total
- 1ml insulin syringes to Reconstitute (Or you can just use the 0.5ml syringes)
- 0.5ml insulin syringes for Daily Dosing.

Mixing & Preparation Instructions

- ◆ Reconstitute Peptides Separately Using Bacteriostatic Water (BAC Water)
 - GHK-Cu 50 mg/mL - Reconstitute with 3mL of BAC Water
 - BPC-157 10 mg/mL - Reconstitute with 2mL of BAC Water
 - TB-500 10 mg/mL - Reconstitute with 1mL of BAC Water
 - Let each peptide sit for a few minutes after adding BAC, gently swirl (do not shake) until fully dissolved.
 - Refrigerate immediately.

Recommended Doses

- ◆ Doseage after reconstituting
 - 2000mcg/2mg GHK-CU (12 units)/daily
 - 600mcg/0.6mg BPC-157 (12 units)/daily
 - 2000-2500 mcg = (20-25 units)/ 3x per week
- Draw BCP-157, TB500, GHK-cu into the same syringe.
- Inject subcutaneously in the lower abdomen or love handle.
- Rotate injection sites.

17 Suggested Monthly/Yearly Cycle Schedule

- 4-6 Weeks On
- 4 Weeks Off

How Much To Purchase for 6 Week

- GHK-cu (2 vials)
- BPC157 (3 vials)
- TB500 (3 vials)
- BAC Water (36mls)
- You will need to order additional BAC water and syringes
 - www.BacteriostaticWater.com: NEWCUST for one time code
 - www.peptidetest.com
 - www.ebac-water.com
 - www.atlanticmedsupply.com

Storage

- Dry lyophilized powder vials can be frozen.
- Store peptide blend in refrigerator (2-8°C) and reconstituting.
- Do not freeze.reconstituted vials.
- Use within 30 days of reconstituting.

Safety Notes

- Always use a new sterile insulin syringe and alcohol swab for each injection.
- Monitor for redness, irritation, or systemic effects.
- Discontinue if unexpected reactions occur.
- Peptides are for research/off-label use and should be monitored by a medical provider.