

If you hit a plateau in your weight loss journey with Semaglutide you can stack the below peptides:

SEMAGLUTIDE 10mg – FOR WEIGHT LOSS

PRECISION PEPTIDE CO. – Code: PEP5 (5% OFF)

<https://precisionpeptideco.com/ref/6/>

- FREE SHIPPING ON ALL ORDERS OVER \$100 (US only)
- FREE 2ml Bacteriostatic Water with every purchase!



WEIGHT LOSS PEPTIDE:

Semaglutide 10mg: <https://precisionpeptideco.com/product/semaglutide-10mg/>

BAC Water: <https://precisionpeptideco.com/product/bacteriostatic-water/>

Code: PEP5

DOSAGE:

The usual adult dose for weight loss is:

- Weeks 1 through 4: 0.25 mg injected subcutaneously once a week
- Weeks 5 through 8: 0.5 mg injected subcutaneously once a week
- Weeks 9 through 12: 1 mg injected subcutaneously once a week
- Weeks 13 through 16: 1.7 mg injected subcutaneously once a week
- Week 14+: 2.4 mg injected subcutaneously once a week

Some people begin to see results within the first eight weeks, but most begin to see results within 12 weeks. Clinical trials have shown that study participants taking a weekly dose of Semaglutide had an average reduction in body weight of 5-10 percent.

Stack (or combine):

1. Tesofensine tablets with Semaglutide (if you plateau) It is recommend to start with 1/2 a pill daily for 2-4 weeks and then go to 1 pill a day (500mcg) <https://precisionpeptideco.com/product/tesofensine-500mcg/>

Possible side effects:

- Constipation • Diarrhea • Dry mouth • Headache • Changes in blood pressure (increase blood pressure) • Increased heart rate • Insomnia • Nausea

2. Cagrilintide 10mg: (This can be combined with Semaglutide if needed)

<https://precisionpeptideco.com/product/cagrilintide-10mg/>

OR instead of Stacking Use:

CagriSem 5-5mg: (If you want to lose less than 30 lbs.)

<https://precisionpeptideco.com/product/cagrisem-5-5mg/>

Discount Code: PEP5

Dosage and Administration: (Dilute 10mg Vial with 2ml BAC Water)

- Weeks 1 through 4: 0.25 mg subcutaneously once a week.
- Weeks 5 through 8: 0.5 mg subcutaneously once a week.
- Weeks 9 through 12: 1 mg subcutaneously once a week.
- Weeks 13 through 16: 1.7 mg subcutaneously once a week.